



# Bt Worm & Caterpillar KILLER

Controls Worms & Caterpillars on Fruits, Vegetables, Ornamentals & Shade Trees

Kills Tomato Hornworms, Tent Caterpillars, Gypsy Moths & Others Listed



#### ACTIVE INGREDIENT:

\**Bacillus thuringiensis* subspecies *kurstaki* strain SA-12solids, spores and Lepidopteran active toxins (At least 6 million viable spores per mg.)\* ..... 15.0%

OTHER INGREDIENTS ..... 85.0%

TOTAL ..... 100.0%

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply **Bt Worm & Caterpillar KILLER** when worms or caterpillars are first noticed, then repeat at five (5) to seven (7) day intervals while they are active. Apply more frequently to control heavy infestations. Apply thoroughly to top and bottom of foliage. Reapply after heavy rains.

**Bt Worm & Caterpillar KILLER** must be eaten by worms or caterpillars to be effective. After ingesting the insecticide, they immediately stop feeding, though they may otherwise appear to be unaffected for several days. Best results are obtained by treatments when worms are small; they must be actively feeding on treated, exposed foliage. You may apply **Bt Worm & Caterpillar KILLER** up to and on the day of harvest.

#### HOW TO APPLY

Always shake or stir **Bt Worm & Caterpillar KILLER** thoroughly before use. Partially fill sprayer with water before adding the application rate amount of product, then mix in product thoroughly and add remaining amount of water.

#### GENERAL INFORMATION

This product is a bacterium which is selectively toxic to many moth and butterfly larvae (Caterpillars). The insects stop feeding and die within 2 - 3 days of ingestion. This product may be applied up to the day of harvest.

**CONTROLS** rehdumped caterpillars, cabbage looper, omnivorous and fruit tree leaf roller.

**USE** on broccoli, cabbage, mustard greens, melons, tomatoes and others as listed.

**APPLY** using hose-end or pressurized sprayers. Can be applied up to day of harvest.

#### HOW TO APPLY

##### (SHADE TREES & ORNAMENTALS)

Use at the rate of 15-25ml / 20L of treatment water to control the following leaf-eating worms: Bagworm, Spring Cankerworm, Fall Cankerworm, Gypsy Moth\*, Tent Caterpillar, Elm Spanworm & Fall Webworm

##### (FRUITS AND VEGETABLES)

Mix 15-25ml / 20L of treatment water for a hand sprayer and apply to all plant foliage. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of: Cabbage looper on broccoli, cabbage, cauliflower, celery, collards, kale, lettuce, melons, mustard greens, tomatoes and turnip greens. Also kills imported cabbage worm on broccoli, cabbage, cauliflower, collards, kale, mustard greens and turnip greens as well as tomato hornworm on tomatoes.



**STARX PTE LTD** Co. Reg. No. : 201526417D  
 111, North Bridge Road #27-01/02, Peninsula Plaza Singapore 179098  
 Tel: +65 8112 5611 Fax: +65 6362 0116 Email: starx@singnet.com.sg  
 Contact Person: Dennis Lim Mobile: +65 9055 1616

Packing Size:  
**1L**