

LIQUID COPPER FUNGICIDE



ACTIVE INGREDIENT:

COPPER OCTANOATE (COPPER SOAP)10.0%
 INERT INGREDIENTS90.0%
TOTAL100.00%

Liquid Copper Fungicide controls a large variety of plant diseases including peach leaf curl, powdery mildew, black spot, rust, anthracnose, fire blight and bacterial leaf spot. Use on vegetables, roses, fruits and turf. Can be used up to the day of harvest.

Where to use - For Roses, Fruits & Vegetables / Ornamentals and Turf / Controls Powdery Mildew, Black Spot and Rust.

Directions for Use SHAKE WELL BEFORE USE.

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops or persists.

If on Skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

Crop	Disease Controlled	Application Rate (10 Ltrs of Treatment Water)	Application Advise
Rose & Ornamental Shrubs	Blackspot, Downy mildew, Gray mold (Botrytis), Leafspots, Powdery mildew, Rust	25ml to 35ml	Begin treatment when disease first appears, and repeat at 7 – 10 days intervals as long as needed.
Sycamore	Anthracnose leaf spot	35ml to 45ml	Make first application just before buds begin to swell, and repeat twice, at 7-day intervals.
Turf	Ascochyta leaf blight, Cercospora leaf spots, Dollar spot	40ml to 50ml	For best control, begin treatment 2 weeks before disease normally appears. Alternatively, begin treatment when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. To reduce Ascochyta leaf blight mow less frequently, only as necessary to maintain recommended height. Water before noon to allow grass to dry. Water thoroughly only as required to avoid moisture stress. Apply Liquid Copper Fungicide when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. In frequently diseased areas, prune adjacent trees and shrubs to reduce turf shading and to improve air movement.
	Rust	40ml to 50ml	To reduce rust mow frequently to reduce rust spore production. Water and fertilize lawn as required to avoid moisture and nutrient stress. Water before noon to allow grass to dry. Apply Liquid Copper Fungicide when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. In frequently diseased areas, prune adjacent trees and shrubs to reduce turf shading and to improve air movement.
Chili, Strawberry, Tomato, Leafy Vegetable	Angular leaf spot, Leaf scorch, Mycosphaerella leaf spot, Phomopsis leaf blight, Powdery mildew, Septoria leaf spots	30ml to 40ml	Spray 1 month after planting (or before flowering on established plants) and twice more at 7 to 10 day intervals.)



STARX PTE LTD UEN No. : 201526417D
 15 Yishun Industrial Street 1 #08-20, Win5, Singapore 768091
 Contact Person: Dennis Lim Mobile: +65 9055 1616
 Email: sales@starx.com.sg



www.starx.com.sg

Packing Size:
1L