

# LIQUID COPPER FUNGICIDE



Can be used on fruit, herbs, vegetables and ornamentals



Controls leaf curl, leaf spots, blights, rust, black spot, downy mildew, powdery mildew, anthracnose and phytophthora stem canker



Can be used up to the day of harvest



**LIQUID COPPER FUNGICIDE** (Active Ingredient: Copper Octanoate) controls a large variety of plant diseases including peach leaf curl, powdery mildew, black spot, rust, anthracnose, fire blight and bacterial leaf spot. Use on vegetables, roses, fruits and turf. Can be used up to the day of harvest.

Crop	Disease Controlled	Application Advise
Chili, Strawberry, Tomato, Leafy Vegetable	Angular leaf spot, Leaf scorch, Mycosphaerella leaf spot, Phomopsis leaf blight, Powdery mildew, Septoria leaf spots	Spray 1 month after planting (or before flowering on established plants) and twice more at 7 to 10 day intervals.
Rose & Ornamental Shrubs	Blackspot, Downy mildew, Gray mold (Botrytis), Leafspots, Powdery mildew, Rust	Begin treatment when disease first appears, and repeat at 7 – 10 day intervals as long as needed.
Sycamore	Anthracnose leaf spot	Make first application just before buds begin to swell, and repeat twice, at 7 day intervals.



Turf	Ascochyta leaf blight, Cercospora leaf spots, Dollar spot	For best control, begin treatment 2 weeks before disease normally appears. Alternatively, begin treatment when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. To reduce Ascochyta leaf blight mow less frequently, only as necessary to maintain recommended height. Water before noon to allow grass to dry. Water thoroughly only as required to avoid moisture stress. Apply Liquid Copper Fungicide when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. In frequently diseased areas, prune adjacent trees and shrubs to reduce turf shading and to improve air movement.
	Rust	To reduce rust mow frequently to reduce rust spore production. Water and fertilize lawn as required to avoid moisture and nutrient stress. Water before noon to allow grass to dry. Apply Liquid Copper Fungicide when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. In frequently diseased areas, prune adjacent trees and shrubs to reduce turf shading and to improve air movement.
<p><b>REMARKS :</b> When to use Liquid Copper Fungicide, do not expect it's to cure an existing fungal disease. The product works by protecting plants against the development of new infections. Ideally, apply Liquid Copper Fungicide before fungus is visible. Otherwise, apply the product immediately when you first notice signs of fungal disease. If the fungus is on fruit trees or vegetable plants, you can safely continue to spray every 7 to 10 days until harvest. If possible, spray plants when you'll have at least 12 hours of dry weather following application.</p>		



**STARX PTE LTD** UEN No. : 201526417D  
 15 Yishun Industrial Street 1 #08-20, Win5, Singapore 768091  
 Contact Person: Dennis Lim Mobile: +65 9055 1616  
 Email: sales@starx.com.sg



www.starx.com.sg

Packing Size:  
**500ML**