



LIME SULFUR

Insects & Disease Control

For Citrus, Fruit Trees, Shrubs, Rose and Ornamentals



CONTROLS

- Mites
- Scale
- Powdery Mildew
- Leaf curl
- Rust

ACTIVE INGREDIENT: Calcium polysulfide99.0%
OTHER INGREDIENTS:1.0%
TOTAL:100.0%

LIME SULPHUR controls a range of fungal diseases and pests on fruit trees, tomatoes, grapes, roses, other ornamentals and edible garden plants. No withholding period.

Pest Control Class: Fungicide, Insecticide, Miticide

Fungus & Disease Control: Anthracnose, Brown Rot, Leaf Spot, Peach Leaf Curl, Powdery Mildew, Rust, Scab, Shothol

Insect Control: Aphids, Mites, Scale, Thrips

FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Call poison control center or doctor immediately for treatment advice.
 - Have person drink a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
 - If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

Plant	Pest / Disease	Application Frequency
Grape Vines	Bud mite, Erinose mite, Rust mite, Vine bunch mite	Apply every two to three weeks (14 days to 21 days).
Citrus	White louse scale, Rust mite, Bud mite, Sooty blotch Citrus rust mite	
Roses & Ornamentals	Rust & Powdery mildew Two spotted mite & Scale insects	
Tomatoes & Vegetables	Powdery mildew & Rust Tomato mites & Two spotted mites	
Stone Fruit (almonds, apricots, peaches, plums, nectarines)	Black spot, Leaf curl, Bryobia mite, Freckle, Rust, Shot hole, Brown rot (blossom blight), San Jose scale, Frosted scale, Powdery mildew	

Directions For Use: SHAKE WELL BEFORE USE.

How to use : DO NOT spray when temperatures exceed 32°C or when soil is dry and plants are suffering from moisture stress. DO NOT apply when freezing weather is expected.

DO NOT apply within 14 days of an oil spray.



STARX PTE LTD UEN No. : 201526417D
 15 Yishun Industrial Street 1 #08-20, Win5, Singapore 768091
 Contact Person: Dennis Lim Mobile: +65 9055 1616
 Email: sales@starx.com.sg



www.starx.com.sg

Packing Size:
500ML