

LIQUID COPPER FUNGICIDE



ACTIVE INGREDIENT

Copper Octanoate (Copper Soap).....	10.0%
Inert Ingredients.....	90.0%
Total	100.00%

Liquid Copper Fungicide controls a large variety of plant diseases including peach leaf curl, powdery mildew, black spot, rust, anthracnose, fire blight and bacterial leaf spot. Use on vegetables, roses, fruits and turf. Can be used up to the day of harvest.

WHERE TO USE

For roses, fruits & vegetables / ornamentals and turf / controls powdery mildew, black spot and rust.

DIRECTIONS FOR USE

Shake well before use

Crop	Disease Controlled	Application Rate (per 10L of Water)	Application Advise
Rose & Ornamental Shrubs	Blackspot, Downy mildew, Gray mold (Botrytis), Leafspots, Powdery mildew, Rust	25ml - 35ml	Apply at first appearance of disease. Repeat every 7-10 days if needed.
Sycamore	Anthrachnose leaf spot	35ml - 45ml	Apply before buds begin to swell and repeat twice every 7 days.
Turf	Ascochyta leaf blight, Cercospora leaf spots, Dollar spot	40ml - 50ml	Begin treatment 2 weeks before disease appears or at first signs; repeat every 7-10 days as needed. Reduce disease by proper mowing, morning watering, and improving air movement.
	Rust	40ml - 50ml	Mow frequently, water before noon, and avoid plant stress. Apply Liquid Copper Fungicide at first signs and repeat every 7-10 days as needed, with good air movement.
Chili, Strawberry, Tomato, Leafy Vegetable	Angular leaf spot, Leaf scorch, Mycosphaerella leaf spot, Phomopsis leaf blight, Powdery mildew, Septoria leaf spots	30ml - 40ml	Spray 1 month after planting (or before flowering) and repeat twice every 7-10 days.



STARX PTE LTD UEN No. : 201526417D

📍 15 Yishun Industrial Street 1 #08-20, Win5, Singapore 768091

☎ +65 8820 7616 / +65 9055 1616

✉ sales@starx.com.sg



www.starx.com.sg

Packing Size:

1L