

All Seasons

Horticultural White Summer Oil



- Dormant & growing season insect spray
- Kills all stages of insects including eggs
- May be used up to the day of harvest

ACTIVE INGREDIENT:

White Summer Oil	72.00 %
Inert Material	28.00 %
TOTAL:	100.00 %

Insects	Crops	Rate Per		Spray Volume per Hectare	Maximum Sprays per Season
		10L of Water	Hectare		
Scale insects (<i>Coccus mangiferae</i> , <i>Aspidiotus orientalis</i>)	Papaya, Citrus, Guava	17 ml	1.5 liter	900 liter	3 times
	Mango	17 ml	1.5 liter		
	Orchid, Rose, Hibiscus, rosasinensis	28 ml	1.7 liter	600 liter	

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in Eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Note to Physician:

This product contains petroleum distillates and may pose an aspiration pneumonia problem.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves when using or handling this product.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Apply this product only as specified on the label.

ROSES, FLOWERING SHRUBS & ORNAMENTALS

SHAKE WELL. Spray to wet entire surface of plant. Give special attention to underside of limbs and leaves.

PLANT	PESTS	TIMING/RESTRICTIONS
Flowering and Foliage Plants including Roses and Other Flowering Shrubs	Aphids, Adelgids, Eriophyid Mites, Gall Mites, Honey Locust Plant Bug, Lace Bug, Leaf Beetle Larvae, Leafminer (non-larvae), Leafrollers, Mealybugs, Psyllids (immature), Red Mites, Sawfly (larvae), Scales (immature), Spider Mites, Webworms, Whiteflies, Blackspot, Powdery Mildew, Young Caterpillars	Summer (Foliar or Cover): Spray as needed.
Foliage Ornamentals & Bedding Plants	Aphids, Adelgids	Summer (Foliar or Cover): Spray as needed. Application safety during bloom period must be determined for each individual species of plant to be treated by conducting a small test.
Non-Fruit-Bearing Trees: Apple, Cherry, Peach, Pear	Mites Powdery Mildew	Apply every 10-14 days. Spray before disease first appears. Apply every 10-14 days.
Shade Trees** Ornamental Trees** Shrubs (including Conifers, Deciduous and Broadleaf Evergreens & Woody Ornamentals)	Aphids, Adelgids, Eriophyid Mites, Gall Mites, Hemlock Woolly Adelgids, Honey Locust Plant Bug, Lace Bug, Leaf Beetle Larvae, Leafminer (non-larvae), Leafrollers, Mealybugs, Psyllids (immature), Red Mites, Sawfly (larvae), Scales (immature), Spider Mites, Webworms, Whiteflies, Blackspot, Powdery Mildew, Young Caterpillars	Summer (Foliar or Cover): Spray as needed. Use enough spray solution to completely penetrate the leaf canopy and cover both top and bottom of all the leaves and stems until wet but without significant runoff. Fall Dormant applications: Limit use only to: American Red Oak, Japanese Black Pine, Dogwood, Weeping Cherry, Cornelian Cherry, Crabapple, Norway Maple, Purple Plum. Restriction: Do not spray when there is obvious moisture deficit in leaves, or the plant is under stress. This product may cause discoloration of Blue Spruce. Do not use on conifers or ferns not tolerant to oil sprays (e.g. Alberta Dwarf Spruce).



STARX PTE LTD UEN No. : 201526417D

15 Yishun Industrial Street 1 #08-20, Win5, Singapore 768091

Contact Person: Dennis Lim Mobile: +65 9055 1616

Email: sales@starx.com.sg



www.starx.com.sg

Packing Size:

1L